

Generating the mapping class group by torsion elements of small order.

Naoyuki Monden

Abstract

We show that the mapping class group of a closed oriented surface of genus at least three is generated by 3 elements of order 3 and by 4 elements of order 4. Note that the mapping class group cannot be generated by finitely many torsion elements of same order if genus is equal to one or two.

1 Introduction

Let Σ_g denote a closed, oriented surface of genus g , and let $\mathcal{M}_g$ denote its mapping class group, which is the group of homotopy classes of orientation-preserving homeomorphisms.

The study of generators of $\mathcal{M}_g$ was pioneered by Dehn. Dehn [De] proved that $\mathcal{M}_g$ is generated by a finite set of Dehn twists. Lickorish [Li] showed that $3g - 1$ Dehn twists generate $\mathcal{M}_g$. This number was improved to $2g + 1$ by Humphries [Hu]. Humphries proved, moreover, that in fact the number $2g + 1$ is minimal; i.e. $\mathcal{M}_g$ cannot be generated by $2g$ (or less) Dehn twists.

It is classical problem to find small generating sets and torsion generating sets for $\mathcal{M}_g$. Maclachlan [Ma] proved that the moduli space is simply connected as a topological space by showing that $\mathcal{M}_g$ is generated by torsion elements. McCarthy and Papadopoulos [MP] proved that $\mathcal{M}_g$ is generated by infinitely many conjugates of a single involution for $g \geq 3$. Luo [Luo] discovered a first finite set of involutions which generate $\mathcal{M}_g$ for $g \geq 3$. Luo posed the question of whether there is a universal upper bound, independent of g , for the number of torsion elements needs to generate $\mathcal{M}_g$. Brendle and Farb answered Luo's question. Brendle and Farb [BF] proved that $\mathcal{M}_g$ is generated by 3 elements of order $2g + 2$, $4g + 2$ and 2 (or g). More, Korkmaz [Ko] showed that $\mathcal{M}_g$ is generated by 2 torsion elements, each of order $4g + 2$. Brendle and Farb [BF] also constructed a generating set of $\mathcal{M}_g$ for $g \geq 3$ consisting of 6 involutions. Kassabov [Ka] improved their method to show that $\mathcal{M}_g$ is generated by 4 involutions if $g \geq 7$, 5 involutions if $g \geq 5$ and 6 involutions if $g \geq 3$.

For all $g \geq 1$, the elements of order 2, 3 and 4 are in $\mathcal{M}_g$. In the present paper, we construct a generating set of $\mathcal{M}_g$ consisting of elements of small order (> 2).

Main Theorem. *For $g \geq 3$, $\mathcal{M}_g$ can be generated by 3 elements of order 3 and by 4 elements of order 4.*

2 Preliminaries

Let c be a simple closed curve on Σ_g . Then the (right hand) Dehn twist T_c about c is the homotopy class of the homeomorphism obtained by cutting Σ_g along c , twisting one of the side by 360° to the right and gluing two sides of c back to each other. Figure 1 shows the Dehn twist about the curve c . We will




Figure 1: The Dehn twist

denote by T_c the Dehn twist about the curve c .

We recall the following lemmas and theorems.

Lemma 1. *For any homeomorphism h of the surface Σ_g , the Dehn twists around the curves c and $h(c)$ are conjugate in the mapping class group $\mathcal{M}_g$,*

$$T_{h(c)} = hT_ch^{-1}.$$

Lemma 2. *Let c and d be two simple closed curves on Σ_g . If c is disjoint from d , then*

$$T_cT_d = T_dT_c$$

Lemma 3. *If the geometric intersection number of c and d is one, then*

$$T_cT_dT_c = T_dT_cT_d$$

Theorem 4 (Lickorish). *We denote the curves $\alpha_i, \beta_i, \gamma_i$ as shown in Figure 2. Then $\mathcal{M}_g$ is generated by $T_{\alpha_1}, \dots, T_{\alpha_g}, T_{\beta_1}, \dots, T_{\beta_g}, T_{\gamma_1}, \dots, T_{\gamma_{g-1}}$.*




Figure 2: Lickorish's generators

We call $T_{\alpha_1}, \dots, T_{\alpha_g}, T_{\beta_1}, \dots, T_{\beta_g}, T_{\gamma_1}, \dots, T_{\gamma_{g-1}}$ Lickorish's generators.

Theorem 5 (Humphries). *$\mathcal{M}_g$ is generated by $T_{\alpha_1}, T_{\alpha_2}, T_{\beta_1}, \dots, T_{\beta_g}, T_{\gamma_1}, \dots, T_{\gamma_{g-1}}$.*

We call $T_{\alpha_1}, T_{\alpha_2}, T_{\beta_1}, \dots, T_{\beta_g}, T_{\gamma_1}, \dots, T_{\gamma_{g-1}}$ Humphries's generators.

3 Generating the mapping class group by 3 elements of order 3

In this section we prove that the mapping class group $\mathcal{M}_g$ is generated by 3 elements of order 3. We assume that $g \geq 3$.

3.1 Construction of elements of order 3

We construct two elements of order 3 by cutting and gluing surfaces. We take the curves x_1, x_2 and the separating curve δ like Figure 3.




Figure 3: the curves x_1, x_2, δ

3.1.1 Odd genus

We assume that g is odd. We construct an element f of order 3.

We cut Σ_g along the curves $\gamma_1, \gamma_2, \alpha_3$ and $\alpha_{2k}, \gamma_{2k}, \alpha_{2k+1}$ ($k = 2, \dots, \frac{g-1}{2}$) to obtain $\frac{g-1}{2}$ surfaces $S_1, S_2, \dots, S_{\frac{g-1}{2}}$ as shown in Figure 4. S_1 is a sphere with $\frac{3(g+1)}{2}$ boundary components and S_k ($k = 2, \dots, \frac{g-1}{2}$) is a pair of pants bounded by $\alpha_{2k}, \gamma_{2k}, \alpha_{2k+1}$.




Figure 4: Cutting the surface

We embed $S_1, S_2, \dots, S_{\frac{g-1}{2}}$ in $\mathbb{R}^3$ so that they are invariant under $\frac{2\pi}{3}$ -rotations $f_1, f_2, \dots, f_{\frac{g-1}{2}}$, respectively (cf. Figure 5).

Since the homeomorphisms $f_1, f_2, \dots, f_{\frac{g-1}{2}}$ coincide on the boundaries, they naturally define a homeomorphism $f : \Sigma_g \rightarrow \Sigma_g$ of order 3. f acts on the curves on Σ_g as follows:

$$\begin{aligned} f^2(\gamma_1) &= f(\gamma_2) = \alpha_3, & f(\beta_2) &= \beta_3, \\ f^2(\alpha_2) &= f(x_2) = x_1, \\ f^2(\alpha_{2k}) &= f(\gamma_{2k}) = \alpha_{2k+1}, & f(\beta_{2k}) &= \beta_{2k+1} \quad (k = 2, \dots, \frac{g-1}{2}). \end{aligned}$$

We construct an element h of order 3.

We cut Σ_g along the curves $\alpha_{2j-1}, \gamma_{2j-1}, \alpha_{2j}$ ($j = 1, \dots, \frac{g-1}{2}$) and δ to obtain $\frac{g+3}{2}$ surfaces $S'_1, S'_2, \dots, S'_{\frac{g+1}{2}}, S'_{\frac{g+3}{2}}$ as shown in Figure 6. S'_1 is a sphere with




Figure 5: $\mathbb{Z}_3$ -symmetry of Σ_g




Figure 6: Cutting the surface, II

$\frac{3g+1}{2}$ boundary components, S'_j ($j = 1, \dots, \frac{g-1}{2}$) is a pair of pants bounded by $\alpha_{2j-1}, \gamma_{2j-1}, \alpha_{2j}$ and $S'_{\frac{g+3}{2}}$ is a torus bounded by δ .

We embed $S'_1, S'_2, \dots, S'_{\frac{g+1}{2}}$ in $\mathbb{R}^3$ so that they are invariant under $\frac{2\pi}{3}$ -rotations $h_1, h_2, \dots, h_{\frac{g+1}{2}}$, respectively (cf. Figure 7). We define that $h_{\frac{g+3}{2}} = (T_{\beta_g} T_{\alpha_g})^2$. Note that $h_1^3 = T_\delta^{-1}$, $h_{\frac{g+3}{2}}^3 = T_\delta$ and $h_{\frac{g+3}{2}}(\alpha_g) = \beta_g$.




Figure 7: $\mathbb{Z}_3$ -symmetry of Σ_g , II

Since the homeomorphisms $h_1, h_2, \dots, h_{\frac{g+1}{2}}, h_{\frac{g+3}{2}}$ coincide on the boundaries, they naturally define a homeomorphism $h : \Sigma_g \rightarrow \Sigma_g$ of order 3. h acts on the curves on Σ_g as follows:

$$\begin{aligned} h(\alpha_g) &= \beta_g, \\ h^2(\alpha_{2j-1}) &= h(\gamma_{2j-1}) = \alpha_{2j}, \quad h(\beta_{2j-1}) = \beta_{2j} \quad (j = 1, \dots, \frac{g-1}{2}). \end{aligned}$$

3.1.2 Even genus

We assume that g is even. By the similar arguments of the case of odd genus we construct f and h which are order 3.

We cut Σ_g along the curves $\gamma_1, \gamma_2, \alpha_3, \delta$ and $\alpha_{2k}, \gamma_{2k}, \alpha_{2k+1}$ ($k = 2, \dots, \frac{g-2}{2}$) to obtain $\frac{g}{2}$ surfaces $S_1, S_2, \dots, S_{\frac{g-2}{2}}, S_{\frac{g}{2}}$ as shown in Figure 8. S_1 is a sphere with $\frac{3g+2}{2}$ boundary components, S_k ($k = 2, \dots, \frac{g-2}{2}$) is a pair of pants bounded by $\alpha_{2k}, \gamma_{2k}, \alpha_{2k+1}$ and $S_{\frac{g}{2}}$ is a torus bounded by δ .

We embed $S_1, S_2, \dots, S_{\frac{g-2}{2}}$ in $\mathbb{R}^3$ so that they are invariant under $\frac{2\pi}{3}$ -rotations $f_1, f_2, \dots, f_{\frac{g-2}{2}}$, respectively (cf. Figure 9). We define that $f_{\frac{g}{2}} = (T_{\beta_g} T_{\alpha_g})^2$.




Figure 8: Cutting the surface, III




Figure 9: $\mathbb{Z}_3$ -symmetry of Σ_g , III

Since the homeomorphisms $f_1, f_2, \dots, f_{\frac{g-2}{2}}, f_{\frac{g}{2}}$ coincide on the boundaries, they naturally define a homeomorphism $f : \Sigma_g \rightarrow \Sigma_g$ of order 3. f acts on the curves on Σ_g as follows:

$$\begin{aligned} f^2(\gamma_1) &= f(\gamma_2) = \alpha_3, & f(\beta_2) &= \beta_3, \\ f^2(\alpha_2) &= f(x_2) = x_1, & f(\alpha_g) &= \beta_g, \\ f^2(\alpha_{2k}) &= f(\gamma_{2k}) = \alpha_{2k+1}, & f(\beta_{2k}) &= \beta_{2k+1} \quad (k = 2, \dots, \frac{g-2}{2}). \end{aligned}$$

We cut Σ_g along the curves $\alpha_{2j-1}, \gamma_{2j-1}, \alpha_{2j}$ ($j = 1, \dots, \frac{g}{2}$) to obtain $\frac{g+2}{2}$ surfaces $S'_1, S'_2, \dots, S'_{\frac{g+2}{2}}$ as shown in Figure 10. S'_1 is a sphere with $\frac{3g}{2}$ boundary components and S'_{j+1} ($j = 1, \dots, \frac{g}{2}$) is a pair of pants bounded by $\alpha_{2j-1}, \gamma_{2j-1}, \alpha_{2j}$.




Figure 10: Cutting the surface, IV

We embed $S'_1, S'_2, \dots, S'_{\frac{g+2}{2}}$ in $\mathbb{R}^3$ so that they are invariant under $\frac{2\pi}{3}$ -rotations $h_1, h_2, \dots, h_{\frac{g+2}{2}}$, respectively (cf. Figure 11).




Figure 11: $\mathbb{Z}_3$ -symmetry of Σ_g , IV

Since the homeomorphisms $h_1, h_2, \dots, h_{\frac{g+2}{2}}$ coincide on the boundaries, they naturally define a homeomorphism $h : \Sigma_g \rightarrow \Sigma_g$ of order 3. h acts on the curves on Σ_g as follows:

$$h^2(\alpha_{2j-1}) = h(\gamma_{2j-1}) = \alpha_{2j}, \quad h(\beta_{2j-1}) = \beta_{2j} \quad (j = 1, \dots, \frac{g}{2}).$$

3.2 Generating the Dehn twist by 3 elements of order 3

We generate the Dehn twist by 3 elements of order 3. The basic idea is to use the *lantern relation* which was discovered by Dehn and rediscovered by Johnson [Jo].




Figure 12: The Lantern Relation

The *lantern relation* is read as follows :

$$T_{\alpha_1} T_{\gamma_1} T_{\gamma_2} T_{\alpha_3} = T_{\alpha_2} T_{x_1} T_{x_2}.$$

where the curves $\alpha_1, \alpha_2, \alpha_3, \gamma_1, \gamma_2, x_1$ and x_2 are shown in Figure 3 and Figure 12. Since $\alpha_1, \gamma_1, \gamma_2$ and α_3 are disjoint each other and α_2, x_1 and x_2 , by Lemma 2 we can rewrite the relation as

$$T_{\alpha_1} = (T_{\alpha_2} T_{\gamma_1}^{-1})(T_{x_1} T_{\alpha_3}^{-1})(T_{x_2} T_{\gamma_2}^{-1}). \quad (1)$$

We can find that $f^2(\alpha_2) = x_1$, $f^2(\gamma_1) = \alpha_3$, $f(\alpha_2) = x_2$ and $f(\gamma_1) = \gamma_2$ from the argument of Section 3.1. By using Lemma 1 we see that

$$\begin{aligned} (T_{x_1} T_{\alpha_3}^{-1}) &= f^2(T_{\alpha_2} T_{\gamma_1}^{-1}) f^{-2} \\ (T_{x_2} T_{\gamma_2}^{-1}) &= f(T_{\alpha_2} T_{\gamma_1}^{-1}) f^{-1}. \end{aligned}$$

Since h maps γ_1 to α_2 , we see that

$$T_{\alpha_2} = h T_{\gamma_1} h^{-1}$$

and

$$T_{\alpha_2} T_{\gamma_1}^{-1} = h T_{\gamma_1} h^{-1} T_{\gamma_1}^{-1} = h(T_{\gamma_1} h^{-1} T_{\gamma_1}^{-1}).$$

Let $\bar{h}$ denote $T_{\gamma_1} h^{-1} T_{\gamma_1}^{-1}$. We can now rewrite (1) as

$$T_{\alpha_1} = (h \bar{h})(f^2 h \bar{h} f^{-2})(f h \bar{h} f^{-1}). \quad (2)$$

and hence T_{α_1} is a product of 3 elements of order 3.

3.3 Proof that 3 elements of order 3 generate

We prove Theorem.

Theorem 6. *If $g \geq 3$, $\mathcal{M}_g$ is generated by f , h and $\bar{h}$.*

Proof. Let G_1 denote the group generated by f , h and $\bar{h}$. By the relation (2), T_{α_1} is in G_1 .

Let α and β be simple closed curves on Σ_g . The symbol $\alpha \xrightarrow{f} \beta$ (resp. $\alpha \xrightarrow{h} \beta$) means that $f(\alpha) = \beta$ (resp. $h(\alpha) = \beta$).

The Figure 13 shows that we can send α_1 to all α_i and γ_i by f and h . Therefore, $T_{\alpha_i}, T_{\gamma_i} \in G_1$ for all i .




Figure 13:

Since h (resp. f) maps α_g to β_g in the case of odd (resp. even) genus, $T_{\beta_g} \in G_1$. As shown on Figure 14, we can find that β_g can be sent to β_i for all i by f and h . These shows that $T_{\beta_i} \in G_1$ for all i .




Figure 14:

Since we show that all Lickorish's generators are in G_1 , G_1 is equal to $\mathcal{M}_g$. $\square$

4 Generating the mapping class group by 4 elements by order 4

In this section we prove that $\mathcal{M}_g$ can be generated by 4 elements of order 4. The key point is to use *chain relation*.

4.1 Construction of elements of order 4

We prepare to construct 3 elements of order 4.

We recall *chain relation*. We say that an ordered set of $c_1, \dots, c_n$ of simple closed curves on Σ_g forms an n -chain if the geometric intersection $(c_k, c_{k+1}) = 1$ for $k = 1, \dots, n-1$ and $(c_k, c_l) = 0$ if $|k-l| \geq 2$. If n is odd, the boundary of a regular neighborhood of any n -chain has two componets d_1 and d_2 .

The *chain relation* is read as follow :

For a given n -chain $c_1, \dots, c_n$, if n is odd we have

$$(T_{c_1} T_{c_2} \cdots T_{c_n})^{n+1} = T_{d_1} T_{d_2}$$

We denote the simple closed curves $\alpha'_1, \dots, \alpha'_g$, s and the separating curves $\delta_1, \dots, \delta_{g-1}$ as shown on Figure 15.




Figure 15: The simple closed curves $\alpha'_1, \dots, \alpha'_g$, s and the separating curves $\delta_1, \dots, \delta_{g-1}$

We note that $\alpha_1 = \alpha'_1$ and $\alpha_g = \alpha'_g$.

For $i = 1, 2, \dots, g-2$ we define

$$\rho_i = (T_{\alpha_{i+2}} T_{\beta_{i+2}} T_{\gamma_{i+1}} T_{\beta_{i+1}} T_{\gamma_i} T_{\beta_i} T_{\alpha'_i})^2.$$

We can find that the boundary components of a regular neighborhood of 7-chain $\alpha_{i+2}, \beta_{i+2}, \gamma_{i+1}, \beta_{i+1}, \gamma_i, \beta_i, \alpha'_i$ are δ_{i+2} and δ_{i-1} like Figure 16. We note that




Figure 16:

δ_0 and δ_g is trivial. By *chain relation*, we have

$$\rho_i^4 = (T_{\alpha_{i+2}} T_{\beta_{i+2}} T_{\gamma_{i+1}} T_{\beta_{i+1}} T_{\gamma_i} T_{\beta_i} T_{\alpha'_i})^8 = T_{\delta_{i+2}} T_{\delta_{i-1}}.$$

For each i we see that ρ_i acts on the curves on Σ_g as follow:

$$\rho_i^3(\alpha_{i+2}) = \rho_i^2(\gamma_{i+1}) = \rho_i(\gamma_i) = \alpha'_i, \quad \rho_i^2(\beta_{i+2}) = \rho_i(\beta_{i+1}) = \beta_i.$$

In paticular we note that $\rho_1(\alpha_2) = x_1$.

We define

$$\rho' = (T_{\gamma_2} T_{\beta_3} T_{\alpha_3} T_s T_{\gamma_1} T_{\beta_1} T_{\alpha'_1})^2.$$

We see that the boundary component of a regular neighborhood of 7-chain $\gamma_2, \beta_3, \alpha_3, s, \gamma_1, \beta_2, \alpha'_1$ in Figure 17 is δ_3 . By *chain relation* we have $(\rho')^4 = T_{\delta_3}$.

ρ' acts on the curves on Σ_g as follow:

$$(\rho')^3(\gamma_2) = (\rho')^2(\alpha_3) = \rho'(\gamma_1) = \alpha'_1, \quad (\rho')^3(x_2) = (\rho')^2(\alpha_2) = \rho'(x_2) = \alpha_2.$$




Figure 17:

In particular we note that $\rho'(\alpha_2) = x_2$.

We define

$$\tau_i = (T_{\alpha'_{i+2}} T_{\beta_{i+2}} T_{\gamma_{i+1}} T_{\beta_{i+1}} T_{\gamma_i} T_{\beta_i} T_{\alpha_i})^2$$

We can find that the boundary components of a regular neighborhood of 7-chain $\alpha'_{i+2}, \beta_{i+2}, \gamma_{i+1}, \beta_{i+1}, \gamma_i, \beta_i, \alpha_i$ are δ_{i+2} and δ_{i-1} like Figure 18. By *chain relation* we have $\tau_i^4 = (T_{\alpha'_{i+2}} T_{\beta_{i+2}} T_{\gamma_{i+1}} T_{\beta_{i+1}} T_{\gamma_i} T_{\beta_i} T_{\alpha_i})^8 = T_{\delta_{i+2}} T_{\delta_{i-1}}$.




Figure 18:

For each i we see that τ_i acts on the curves on Σ_g as follow:

$$\tau^3(\alpha'_{i+2}) = \tau^2(\gamma_{i+1}) = \tau(\gamma_i) = \alpha_i \quad \tau^2(\beta_{i+2}) = \tau(\beta_{i+1}) = \beta_i.$$

For $j = 1, 2, \dots, g$ we define

$$\sigma_j = T_{\alpha_j} T_{\beta_j} T_{\alpha'_j}.$$

We can find that the boundary components of a regular neighborhood of 3-chain $\alpha_j, \beta_j, \alpha'_j$ like Figure 19 is δ_j and δ_{j-1} . By Lemma 3 and *chain relation* we have $\tau_j^4 = T_{\delta_j} T_{\delta_{j-1}}$.




Figure 19:

For each j we see that $\sigma_j(\alpha_j) = T_{\alpha_j} T_{\beta_j} T_{\alpha'_j}(\alpha_j) = \beta_j$.

4.1.1 The genus is $3m$

We assume that $g = 3m$. We construct an element ϕ of order 4. We define

$$\phi = \begin{cases} \rho_{g-2}^{-1} \rho_{g-5} \cdots \rho_7^{-1} \rho_4 \rho_1^{-1} & (m(=\frac{g}{3}) \text{ is odd}) \\ \rho_{g-2} \rho_{g-5}^{-1} \cdots \rho_7^{-1} \rho_4 \rho_1^{-1} & (m \text{ is even}). \end{cases}$$

We can find that $\phi^4 = T_{\delta_{g-3}}^{\mp 1} (T_{\delta_{g-3}} T_{\delta_{g-6}})^{\pm 1} \cdots (T_{\delta_9} T_{\delta_6})^{-1} (T_{\delta_6} T_{\delta_3}) T_{\delta_3}^{-1} = 1$.
 ϕ acts on the curves on Σ_g as follow:

$$\begin{aligned} \alpha_2 &= \phi(x_1), & \phi^3(\alpha'_1) &= \phi^2(\gamma_1) = \phi(\gamma_2) = \alpha_3, & \phi^2(\beta_1) &= \phi(\beta_2) = \beta_3 \\ \alpha'_4 &= \phi(\gamma_4) = \phi^2(\gamma_5) = \phi^3(\alpha_6), & \beta_4 &= \phi(\beta_5) = \phi^2(\beta_6) \\ \vdots & & \vdots & \\ \phi^3(\alpha'_{g-2}) &= \phi^2(\gamma_{g-2}) = \phi(\gamma_{g-1}) = \alpha_g, & \phi^2(\beta_{g-2}) &= \phi(\beta_{g-1}) = \beta_g & (m \text{ is odd}), \\ (\alpha'_{g-2} = \phi(\gamma_{g-2}) = \phi^2(\gamma_{g-1}) = \phi^3(\alpha_g), & \beta_{g-2} = \phi(\beta_{g-1}) = \phi^2(\beta_g) & (m \text{ is even})). \end{aligned}$$

We construct an element ψ of order 4. We define

$$\psi = \begin{cases} \rho_{g-2}^{-1} \rho_{g-5} \cdots \rho_7^{-1} \rho_4 \rho'^{-1} & (m(=\frac{g}{3}) \text{ is odd}) \\ \rho_{g-2} \rho_{g-5}^{-1} \cdots \rho_7^{-1} \rho_4 \rho'^{-1} & (m \text{ is even}). \end{cases}$$

We can find that $\psi^4 = 1$. Since $\rho'(\alpha_2) = x_2$ and $\rho'(\gamma_2) = \alpha_3$, we can easily find that $\psi^{-1}(\alpha_2) = x_2$ and $\psi^{-1}(\gamma_2) = \alpha_3$.

We construct an element ω of order 4. We define

$$\omega = \begin{cases} \sigma_g^{-1} \sigma_{g-1} \tau_{g-5}^{-1} \tau_{g-8} \cdots \tau_8 \tau_5^{-1} \tau_2 \sigma_1^{-1} & (m(=\frac{g}{3}) \text{ is odd}) \\ \sigma_g \sigma_{g-1}^{-1} \tau_{g-5} \tau_{g-8}^{-1} \cdots \tau_8 \tau_5^{-1} \tau_2 \sigma_1^{-1} & (m \text{ is even}). \end{cases}$$

We can find that $\omega^4 = 1$.

ω acts on the curves on Σ_g as follow:

$$\begin{aligned} \alpha_1 &= \omega(\beta_1), & \omega^3(\alpha_2) &= \omega^2(\gamma_2) = \omega(\gamma_3) = \alpha'_4, & \omega^2(\beta_2) &= \omega(\beta_3) = \beta_4, \\ \alpha_5 &= \omega(\gamma_5) = \omega^2(\gamma_6) = \omega^3(\alpha'_7), & \beta_5 &= \omega(\beta_6) = \omega^2(\beta_7), \\ \vdots & & \vdots & \\ \omega^3(\alpha_{g-4}) &= \omega^2(\gamma_{g-4}) = \omega(\gamma_{g-3}) = \alpha'_{g-2}, & \omega^2(\beta_{g-4}) &= \omega(\beta_{g-3}) = \beta_{g-2} & (m \text{ is odd}), \\ (\alpha_{g-4} = \omega(\gamma_{g-4}) = \omega^2(\gamma_{g-3}) = \omega^3(\alpha'_{g-2}), & \beta_{g-4} = \omega(\beta_{g-3}) = \omega^2(\beta_{g-2}) & (m \text{ is even})). \end{aligned}$$

4.1.2 The genus is $3m + 1$

We assume that $g = 3m + 1$.

We define

$$\phi = \begin{cases} \sigma_g \rho_{g-3}^{-1} \rho_{g-6} \cdots \rho_7^{-1} \rho_4 \rho_1^{-1} & (m(=\frac{g-1}{3}) \text{ is odd}) \\ \sigma_g^{-1} \rho_{g-3} \rho_{g-6}^{-1} \cdots \rho_7^{-1} \rho_4 \rho_1^{-1} & (m \text{ is even}). \end{cases}$$

ϕ acts on the curves on Σ_g as follow:

$$\begin{aligned}
& \alpha_2 = \phi(x_1), & \phi^2(\beta_1) = \phi(\beta_2) = \beta_3, \\
& \phi^3(\alpha'_1) = \phi^2(\gamma_1) = \phi(\gamma_2) = \alpha_3, & \beta_4 = \phi(\beta_5) = \phi^2(\beta_6) \\
& \alpha'_4 = \phi(\gamma_4) = \phi^2(\gamma_5) = \alpha_6, & \\
& \vdots & \vdots \\
& \phi^3(\alpha'_{g-3}) = \phi^2(\gamma_{g-3}) = \phi(\gamma_{g-2}) = \alpha_{g-1}, & \phi^2(\beta_{g-3}) = \phi(\beta_{g-2}) = \beta_{g-1} \quad (m \text{ is odd}), \\
& (\alpha'_{g-3} = \phi(\gamma_{g-3}) = \phi^2(\gamma_{g-2}) = \phi^3(\alpha_{g-1}), & \beta_{g-3} = \phi(\beta_{g-2}) = \phi^2(\beta_{g-1}) \quad (m \text{ is even}).
\end{aligned}$$

We define

$$\psi = \begin{cases} \rho_{g-2}^{-1} \rho_{g-5} \cdots \rho_7^{-1} \rho_4 \rho'^{-1} & (m(=\frac{g-1}{3}) \text{ is odd}) \\ \rho_{g-2} \rho_{g-5}^{-1} \cdots \rho_7^{-1} \rho_4 \rho'^{-1} & (m \text{ is even}). \end{cases}$$

We note that $\psi^{-1}(\alpha_2) = x_2$ and $\psi^{-1}(\gamma_2) = \alpha_3$.

We define

$$\omega = \begin{cases} \tau_{g-2}^{-1} \tau_{g-5} \cdots \tau_8 \tau_5^{-1} \tau_2 \sigma_1^{-1} & (m(=\frac{g-1}{3}) \text{ is odd}) \\ \tau_{g-2} \tau_{g-5}^{-1} \cdots \tau_8 \tau_5^{-1} \tau_2 \sigma_1^{-1} & (m \text{ is even}). \end{cases}$$

ω acts on the curves on Σ_g as follow:

$$\begin{aligned}
& \alpha_1 = \omega(\beta_1), & \omega^2(\beta_2) = \omega(\beta_3) = \beta_4, \\
& \omega^3(\alpha_2) = \omega^2(\gamma_2) = \omega(\gamma_3) = \alpha'_4, & \beta_5 = \omega(\beta_6) = \omega^2(\beta_7), \\
& \alpha_5 = \omega(\gamma_5) = \omega^2(\gamma_6) = \omega^3(\alpha'_7), & \\
& \vdots & \vdots \\
& \omega^3(\alpha_{g-2}) = \omega^2(\gamma_{g-2}) = \omega(\gamma_{g-1}) = \alpha'_g, & \omega^2(\beta_{g-2}) = \omega(\beta_{g-1}) = \beta_g \quad (m \text{ is odd}), \\
& (\alpha_{g-2} = \omega(\gamma_{g-2}) = \omega^2(\gamma_{g-1}) = \alpha'_g, & \omega^2(\beta_{g-2}) = \omega(\beta_{g-1}) = \beta_g \quad (m \text{ is even})).
\end{aligned}$$

4.1.3 The genus is $3m + 2$

We assume that $g = 3m + 2$ ($m \geq 2$).

We define

$$\phi = \begin{cases} \sigma_g^{-1} \sigma_{g-1} \rho_{g-4}^{-1} \rho_{g-7} \cdots \rho_7^{-1} \rho_4 \rho_1^{-1} & (m(=\frac{g-2}{3}) \text{ is odd}) \\ \sigma_g \sigma_{g-1}^{-1} \rho_{g-4} \rho_{g-7}^{-1} \cdots \rho_7^{-1} \rho_4 \rho_1^{-1} & (m \text{ is even}). \end{cases}$$

ϕ acts on the curves on Σ_g as follow:

$$\begin{aligned}
& \alpha_2 = \phi(x_1), & \phi^2(\beta_1) = \phi(\beta_2) = \beta_3, \\
& \phi^3(\alpha'_1) = \phi^2(\gamma_1) = \phi(\gamma_2) = \alpha_3, & \beta_4 = \phi(\beta_5) = \phi^2(\beta_6), \\
& \alpha'_4 = \phi(\gamma_4) = \phi^2(\gamma_5) = \phi^3(\alpha_6), & \\
& \vdots & \vdots \\
& \phi^3(\alpha'_{g-4}) = \phi^2(\gamma_{g-4}) = \phi(\gamma_{g-3}) = \alpha_{g-2}, & \phi^2(\beta_{g-4}) = \phi(\beta_{g-3}) = \beta_{g-2} \quad (m \text{ is odd}), \\
& (\alpha'_{g-4} = \phi(\gamma_{g-4}) = \phi^2(\gamma_{g-3}) = \phi^3(\alpha_{g-2}), & \beta_{g-4} = \phi(\beta_{g-3}) = \phi^2(\beta_{g-2}) \quad (m \text{ is even})).
\end{aligned}$$

We define

$$\psi = \begin{cases} \rho_{g-2}^{-1} \rho_{g-5} \cdots \rho_9^{-1} \rho_6 \sigma_5^{-1} \sigma_4 \rho'^{-1} & (m(=\frac{g-2}{3}) \text{ is odd}) \\ \rho_{g-2} \rho_{g-5}^{-1} \cdots \rho_9^{-1} \rho_6 \sigma_5^{-1} \sigma_4 \rho'^{-1} & (m \text{ is even}). \end{cases}$$

We note that ψ acts on the curves on Σ_g as follow:

$$\begin{aligned} \alpha_2 &= \psi(x_2), & \gamma_2 &= \psi(\alpha_3), \\ \psi^3(\alpha'_{g-2}) &= \psi^2(\gamma_{g-2}) = \psi(\gamma_{g-1}) = \alpha_g, & \psi^2(\beta_{g-2}) &= \psi(\beta_{g-1}) = \beta_g \quad (m \text{ is odd}), \\ (\alpha'_{g-2} &= \psi(\gamma_{g-2}) = \psi^2(\gamma_{g-1}) = \psi^3(\alpha_g), & \beta_{g-2} &= \psi(\beta_{g-1}) = \psi^2(\beta_g) \quad (m \text{ is even}). \end{aligned}$$

We define

$$\omega = \begin{cases} \sigma_g \tau_{g-3}^{-1} \tau_{g-6} \cdots \tau_8 \tau_5^{-1} \tau_2 \sigma_1^{-1} & (m(=\frac{g-2}{3}) \text{ is odd}) \\ \sigma_g^{-1} \tau_{g-3} \tau_{g-6}^{-1} \cdots \tau_8 \tau_5^{-1} \tau_2 \sigma_1^{-1} & (m \text{ is even}). \end{cases}$$

ω acts on the curves on Σ_g as follow:

$$\begin{aligned} \alpha_1 &= \omega(\beta_1), & \omega^2(\beta_2) &= \omega(\beta_3) = \beta_4, \\ \omega^3(\alpha_2) &= \omega^2(\gamma_2) = \omega(\gamma_3) = \alpha'_4, & \beta_5 &= \omega(\beta_6) = \omega^2(\beta_7), \\ \alpha'_5 &= \omega(\gamma_5) = \omega^2(\gamma_6) = \omega^3(\alpha'_7), & & \\ \vdots & & \vdots & \\ \omega^3(\alpha_{g-3}) &= \omega^2(\gamma_{g-3}) = \omega(\gamma_{g-2}) = \alpha'_{g-1}, & \omega^2(\beta_{g-3}) &= \omega(\beta_{g-2}) = \beta_{g-1} \quad (m \text{ is odd}), \\ (\alpha_{g-3} &= \omega(\gamma_{g-3}) = \omega^2(\gamma_{g-2}) = \omega^3(\alpha'_{g-1}), & \beta_{g-3} &= \omega(\beta_{g-2}) = \omega^2(\beta_{g-1}) \quad (m \text{ is even}). \end{aligned}$$

4.1.4 The genus is 5

We assume that $g = 5$.

We define

$$\phi = (T_{\alpha_5} T_{\beta_5} T_{\gamma_4}) (T_{\gamma_3} T_{\beta_3} T_{\gamma_2} T_{\beta_2} T_{\gamma_1} T_{\beta_1} T_{\alpha'_1})^{-2}.$$

By *chain reltion*, $(T_{\alpha_5} T_{\beta_5} T_{\gamma_4})^4 = (T_{\gamma_3} T_{\beta_3} T_{\gamma_2} T_{\beta_2} T_{\gamma_1} T_{\beta_1} T_{\alpha'_1})^8 = T_{\alpha_4} T_{\alpha'_4}$ Therefore, we can find that $\phi^4 = 1$.

ϕ acts on the curves on Σ_g as follow :

$$\begin{aligned} \phi^3(\alpha'_1) &= \phi^2(\gamma_1) = \phi(\gamma_2) = \alpha_3, & \phi^2(\beta_1) &= \phi(\beta_2) = \beta_3, \\ \alpha_2 &= \phi(x_1), & \phi(\gamma_4) &= \beta_5. \end{aligned}$$

We define

$$\psi = \sigma_5 \sigma_4 \rho'^{-1}.$$

We can find that $\psi^4 = 1$.

ψ acts on the curves on Σ_g as follow :

$$\psi^3(\alpha'_1) = \psi^2(\gamma_1) = \psi(\alpha_3) = \gamma_2, \quad \psi(\alpha_2) = x_2.$$




Figure 20: the curves s_1 and s_2 .

Let s_1, s_2 the non separating simple closed curves as shown in Figure 20
We define

$$\omega = (T_{\alpha_1} T_{\beta_1} T_{s_1})(T_{\gamma_4} T_{\beta_4} T_{\gamma_3} T_{\beta_3} T_{\gamma_2} T_{\beta_2} T_{\alpha_2})^{-2}.$$

By the *chain relation*, $(T_{\alpha_1} T_{\beta_1} T_{s_1})^4 = (T_{\gamma_4} T_{\beta_4} T_{\gamma_3} T_{\beta_3} T_{\gamma_2} T_{\beta_2} T_{\alpha_2})^8 = T_{\alpha_g} T_{s_2}$. Therefore, we can find that $\omega^4 = 1$.

ω acts on the curves on Σ_g as follow :

$$\omega^3(\alpha_2) = \omega^2(\gamma_2) = \omega(\gamma_3) = \gamma_4, \quad \omega^2(\beta_2) = \omega(\beta_3) = \beta_4.$$

4.2 Generating the Dehn twist by 4 elements of order 4

By using the lantern relation we generate the Dehn twist by 4 elements of order 4.

Since ω maps α_2 to γ_2 , we see that

$$T_{\alpha_2} = \omega T_{\gamma_2} \omega^{-1}$$

and

$$T_{\alpha_2} T_{\gamma_2}^{-1} = \omega T_{\gamma_2} \omega^{-1} T_{\gamma_2}^{-1} = \omega (T_{\gamma_2} \omega^{-1} T_{\gamma_2}^{-1}).$$

Let $\tilde{\omega}$ denote $T_{\gamma_2} \omega^{-1} T_{\gamma_2}^{-1}$. Then we see that $T_{\alpha_2} T_{\gamma_2}^{-1} = \omega \tilde{\omega}$.

In the case of $g \neq 5$, we rewrite the lantern relation as follow :

$$T_{\alpha_1} = T_{\alpha'_1} = (T_{\alpha_2} T_{\gamma_2}^{-1})(T_{x_1} T_{\gamma_1}^{-1})(T_{x_2} T_{\alpha_3}^{-1}). \quad (3)$$

From the argument of Section 4.1.1, 4.1.2 and 4.1.3 we can find that $\phi^{-1}(\alpha_2) = x_1$, $\phi^{-1}(\gamma_2) = \gamma_1$, $\psi^{-1}(\alpha_2) = x_2$ and $\psi^{-1}(\gamma_2) = \alpha_3$. By Lemma 1 show we see that :

$$\begin{aligned} (T_{x_1} T_{\gamma_2}^{-1}) &= \phi^{-1}(T_{\alpha_2} T_{\gamma_1}^{-1}) \phi \\ (T_{x_2} T_{\alpha_3}^{-1}) &= \psi^{-1}(T_{\alpha_2} T_{\gamma_1}^{-1}) \psi. \end{aligned}$$

We can rewrite (3) as

$$T_{\alpha_1} = (\omega \tilde{\omega})(\phi^{-1} \omega \tilde{\omega} \phi)(\psi^{-1} \omega \tilde{\omega} \psi).$$

In the case of $g = 5$, we rewrite the lantern relation as follow :

$$T_{\alpha_3} = (T_{\alpha_2} T_{\gamma_2}^{-1})(T_{x_1} T_{\gamma_1}^{-1})(T_{x_2} T_{\alpha_1}^{-1}). \quad (4)$$

From the argument of Section 4.1.3 we can find that $\phi^{-1}(\alpha_2) = x_1$, $\phi^{-1}(\gamma_2) = \gamma_1$, $\psi(\alpha_2) = x_2$ and $\psi(\gamma_2) = \alpha'_1 = \alpha_1$. By Lemma 1 we see that :

$$\begin{aligned}(T_{x_1} T_{\gamma_1}^{-1}) &= \phi^{-1}(T_{\alpha_2} T_{\gamma_2}^{-1}) \phi \\ (T_{x_2} T_{\alpha_1}^{-1}) &= \psi(T_{\alpha_2} T_{\gamma_2}^{-1}) \psi^{-1}.\end{aligned}$$

We can rewrite (4)

$$T_{\alpha_3} = (\omega \tilde{\omega})(\phi^{-1} \omega \tilde{\omega} \phi)(\psi \omega \tilde{\omega} \psi^{-1}).$$

Hence Dehn twist is a product of 4 elements of order 4.

4.3 Proof that 4 elements of order 4 generate

We prove Theorem.

Theorem 7. *If g is at least 3, $\mathcal{M}_g$ can be generated by ϕ , ψ , ω and $\tilde{\omega}$.*

Proof. Let G_2 denote the group generated by ϕ , ψ , ω and $\tilde{\omega}$.

Let α and β be simple closed curves on Σ_g . The symbol $\alpha \xleftrightarrow{\phi} \beta$ (resp. $\alpha \xleftrightarrow{\psi} \beta$, $\alpha \xleftrightarrow{\omega} \beta$) means that either $\phi(\alpha) = \beta$ or $\phi^{-1}(\alpha) = \beta$ (resp. either $\psi(\alpha) = \beta$ or $\psi^{-1}(\alpha) = \beta$, either $\omega(\alpha) = \beta$ or $\omega^{-1}(\alpha) = \beta$).

If $g \neq 5$, we can find that T_{α_1} is in G_2 . In the case of $g = 3m$ and $g = 3m + 1$, ϕ and ω can map α_1 to all β_i and γ_i as shown in Figure 21. If $g = 3m + 2$, we




Figure 21:

need ψ other than ϕ , ω like Figure 22. Therefore, for all i , T_{β_i} , T_{γ_i} , are in G_2 . Because $\omega(\alpha_2) = \gamma_2$, we can find that $T_{\alpha_2} \in G_2$. Therefore, all Humphries's generators are in G_2 .




Figure 22:

If $g = 5$, T_{α_3} is in G_2 . The Figure 23 shows that we can send α_3 to all Humphries's generators by ϕ , ψ and ω .




Figure 23:

We prove that G_2 is equal to $\mathcal{M}_g$ for $g \geq 3$. □

5 Remarks

5.1 Low genus

In the case of $g = 1, 2$, we note that $\mathcal{M}_g$ can not be generated by elements of same order. By using the argument of MacCarthy and Papadopoulos [MP] and the work of Hirose [Hi], we can see the proof. Hirose listed the Dehn twist presentation of finite order elements for closed oriented surfaces of genera up to 4. We introduce the presentation of finite order elements in the case of $g = 1, 2$. The list is as follow :

genus	elements	order
1	$T_{\beta_1} T_{\alpha_1}$	6
	$T_{\alpha_1} T_{\beta_1} T_{\alpha_1}$	4
2	$T_{\beta_2} T_{\gamma_1} T_{\beta_1} T_{\alpha_1}$	10
	$T_{\beta_2} T_{\beta_2} T_{\gamma_1} T_{\beta_1} T_{\alpha_1}$	8
	$T_{\alpha_2} T_{\beta_2} T_{\gamma_1} T_{\beta_1} T_{\alpha_1}$	6
	$(T_{\alpha_1} T_{\beta_1} T_{\gamma_1} T_{\beta_2} T_{\alpha_2})(T_{\alpha_2} T_{\beta_2} T_{\gamma_1} T_{\beta_1} T_{\alpha_1})^3$	6

MacCarthy and Papadopoulos proved that $\mathcal{M}_2$ can not be generated by elements of order 2. The argument of MacCarthy and Papadopoulos is as follow:

Proof. Let c be a nonseparating simple closed curve and p be the abelianization map given by Powell's result [Po]:

$$\begin{array}{ccc} p : \mathcal{M}_2 & \longrightarrow & \mathbb{Z}_{10} \\ \Psi & & \Psi \\ T_c & \longmapsto & 1 \end{array}$$

We can find that

$$\begin{aligned} p((T_{\beta_2}T_{\gamma_1}T_{\beta_1}T_{\alpha_1})^5) &= p((T_{\beta_2}T_{\beta_2}T_{\gamma_1}T_{\beta_1}T_{\alpha_1})^4) \\ &= p(\{(T_{\alpha_1}T_{\beta_1}T_{\gamma_1}T_{\beta_2}T_{\alpha_2})(T_{\alpha_2}T_{\beta_2}T_{\gamma_1}T_{\beta_1}T_{\alpha_1})^3\}^3) \\ &= 0 \\ p((T_{\alpha_2}T_{\beta_2}T_{\gamma_1}T_{\beta_1}T_{\alpha_1})^3) &= 5. \end{aligned}$$

It is easily to see that $\mathbb{Z}_{10}$ can not be generated by 0 and 5. Therefore we can see that $\mathcal{M}_2$ can not be generated by elements of order 2. $\square$

By the similar proof, we can see that $\mathcal{M}_1$ and $\mathcal{M}_2$ can not be generated by elements of same order.

Remark 8. $\mathcal{M}_1$ and $\mathcal{M}_2$ can be generated by elements of different order. For example, $\mathcal{M}_1$ can be generated by $T_{\beta_1}T_{\alpha_1}$ and $T_{\alpha_1}T_{\beta_1}T_{\alpha_1}$, and $\mathcal{M}_2$ can be generated by $T_{\beta_2}T_{\gamma_1}T_{\beta_1}T_{\alpha_1}$ and $T_{\alpha_2}T_{\beta_2}T_{\gamma_1}T_{\beta_1}T_{\alpha_1}$.

5.2 Lower bound

The order of $\mathcal{M}_g$ is not finite. Therefore, we can easily find that lower bound of the number of generators whose order are 3 (resp. 4) is 2. The author has the following question :

Question. *What is the minimal number of elements of order 3 (resp. 4) required to generate $\mathcal{M}_g$?*

Reference

- [BF] T. E. Brendle and B. Farb. *Every mapping class group is generated by 3 torsion elements and by 6 involutions.* J. Algebra **278** (2004), 187–198.
- [De] M. Dehn. *Papers on group theory and topology.* Springer-Verlag, New York, 1987 (Die Gruppe der Abbildungsklassen, Acta Math. Vol. **69** (1938), 135–206)
- [Hi] S. Hirose. *Presentations of periodic maps on oriented closed surfaces of genera up to 4.* Osaka Journal of Mathematics, to appear.
- [Hu] S. P. Humphries. *Generators for the mapping class group.* Topology of low-dimensional manifolds Proc. Second Sussex Conf. Chelwood Gate 1977, Lecture Notes in Math. **722** (1979), Springer, 44–47.

- [Jo] D. Johnson. *The structure of Torelli group I: A finite set of generators for $\mathcal{I}$* . Ann. of Math. **118** (1983), 423–442.
- [Ka] M. Kassabov. *Generating Mapping Class Groups by Involutions*. arXiv:math.GT/0311455 v1 25 Nov 2003.
- [Ko] M. Korkmaz. *Generating the surface mapping class group by two elements*. Trans. Amer. Math. Soc. **357** (2005), 3299–3310.
- [Li] W. B. R. Lickorish. *A finite set of generators for the homeotopy group of a 2-manifold*. Proc. Cambridge Philos. Soc. **60** (1964), 769–778.
- [Luo] F. Luo. *Torsion Elements in the Mapping Class Group of a Surface*. arXiv:math.GT/0004048 v1 8 Apr 2000.
- [Ma] C. Maclachlan. *Modulus space is simply-connected*. Proc. Amer. Math. Soc. **29** (1971), 85–86.
- [MP] J. MacCarthy and A. Papadopoulos. *Involutions in surface mapping class groups*. Enseign. Math. **33** (1987), 275–290.
- [Po] J. Powell. *Two theorems on the mapping class group of a surface*. Proc. Amer. Math. Soc. **68** (1978), no. 3, 347–350.

Department of Mathematics, Graduate School of Science, Osaka University,
Toyonaka, Osaka 560-0043, Japan

E-mail adress: **n-monden@cr.math.sci.osaka-u.ac.jp**

$$\alpha \xrightarrow{\bar{f}} \beta$$